

SUPPLEMENTARY MATERIAL  
Admissible, Similar Tests:  
A characterization.  
Appendix A.

## APPENDIX A: FINITE-SAMPLE THEORY

A.1. Lemma 1: Weak\* compactness of  $\mathcal{C}(\alpha\text{-s})$ 

DESCRIPTION: The first lemma of this appendix shows that the set of  $\alpha$ -similar tests, denoted  $\mathcal{C}(\alpha\text{-s})$ , is compact relative to the space of essentially bounded measurable functions endowed with the weak\* topology.

RELEVANCE OF LEMMA 1: This lemma will be used to prove part ii) of Theorem 1. The weak\* compactness of  $\mathcal{C}(\alpha\text{-s})$  will allow the application of an essentially complete class Theorem [See Theorem 3, p. 87, Chapter 2 in [Ferguson \(1967\)](#)].

NOTATION: Let  $\mathcal{B}(\mathbb{R}^n)$  denote the Borel  $\sigma$ -algebra on  $\mathbb{R}^n$ . For any set  $\mathcal{S} \in \mathcal{B}(\mathbb{R}^n)$ , let  $\mathcal{B}(\mathbb{R}^n)_{\mathcal{S}}$  denote the sub-space  $\sigma$ -algebra. *Measurability* of the function  $f : \mathcal{S} \rightarrow \mathbb{R}$  is always relative to the measurable spaces  $(\mathcal{S}, \mathcal{B}(\mathbb{R}^n)_{\mathcal{S}})$ - $(\mathbb{R}, \mathcal{B}(\mathbb{R}))$ . The integral of  $f$  with respect to the Lebesgue measure in  $\mathbb{R}^n$  is denoted by  $\int_{\mathcal{S}} f(s) ds$ . *Integration* with respect to a different measure  $\mu$  is denoted  $\int_{\mathcal{S}} f(s) d\mu(s)$  or  $\int_{\mathcal{S}} f d\mu$  if no ambiguity arises. All vectors are column vectors. For notational convenience,  $(a, b)$  will sometimes replace  $(a', b)'$ . The dimension of the column vector “a” is denoted  $d_a$ .

**PRELIMINARIES 1** ( $L^1$  and  $L^\infty$ ): Since the sample space  $\mathbf{X} \in \mathcal{B}(\mathbb{R}^s)$ , the triplet  $(\mathbf{X}, \mathcal{B}(\mathbb{R}^s)_{\mathbf{X}}, \lambda^s)$  is a well-defined  $\sigma$ -finite measure space, where  $\lambda^s$  denotes the Lebesgue measure in  $\mathbb{R}^s$  restricted to  $\mathbf{X}$ . Note that  $\mathcal{B}(\mathbb{R}^s)_{\mathbf{X}} = \mathcal{B}(\mathbf{X})$  whenever  $\mathbf{X}$  is endowed with the sub-space topology relative to  $\mathbb{R}^s$ . Following [Rudin \(2006\)](#), p. 65, let  $L^1(\mathbf{X}, \mathcal{B}(\mathbf{X}), \lambda^s)$  denote the space of all real-valued measurable functions  $f$  that satisfy  $\|f\|_1 \equiv \int_{\mathbf{X}} |f(x)| dx < \infty$ . Let  $L^\infty(\mathbf{X}, \mathcal{B}(\mathbf{X}), \lambda^s)$  denote the class of all essentially bounded real-valued measurable functions ([Rudin \(2006\)](#) p. 66).

REMARK 3: Identify the class of all tests  $\mathcal{C}$  as a subset of  $L^\infty(\mathbf{X}, \mathcal{B}(\mathbf{X}), \lambda^s)$

$$\mathcal{C} \equiv \{ \phi \in L^\infty(\mathbf{X}, \mathcal{B}(\mathbf{X}), \lambda^s) \mid \phi(x) \in [0, 1] \text{ for } \lambda^s\text{-a.e. } x \in \mathbf{X} \}.$$

And note that the elements of any statistical model  $\{f(x; \theta)\}_{\theta \in \Theta}$  are elements of  $L^1(\mathbf{X}, \mathcal{B}(\mathbf{X}), \lambda^s)$ , by the definition of probability density function  $\int_{\mathbf{X}} f(x; \theta) dx = 1 < \infty$  for all  $\theta \in \Theta$ .

**PRELIMINARIES 2** (The dual space of  $L^1$ ): Let  $[L^1(\mathbf{X}, \mathcal{B}(\mathbf{X}), \lambda^s)]^*$  denote the dual space of  $L^1(\mathbf{X}, \mathcal{B}(\mathbf{X}), \lambda^s)$ , i.e., the space of all continuous (w.r.t.  $\|f\|_1$ ) linear functionals on  $L^1(\mathbf{X}, \mathcal{B}(\mathbf{X}), \lambda^s)$ ; see [Rudin \(2005\)](#), p. 56. Let  $\Lambda$  denote an element of the dual space  $[L^1(\mathbf{X}, \mathcal{B}(\mathbf{X}), \lambda^s)]^*$ . By Theorem 6.16 in [Rudin \(2006\)](#), p. 127 and Theorem 1.18 in [Rudin \(2005\)](#), p. 15; the space  $[L^1(\mathbf{X}, \mathcal{B}(\mathbf{X}), \lambda^s)]^*$  is isometrically isomorphic to  $L^\infty(\mathbf{X}, \mathcal{B}(\mathbf{X}), \lambda^s)$ . Therefore, one can identify each functional  $\Lambda$  with a unique element (up to equivalence)  $g \in L^\infty(\mathbf{X}, \mathcal{B}(\mathbf{X}), \lambda^s)$ , and vice versa: for  $f \in L^1(\mathbf{X}, \mathcal{B}(\mathbf{X}), \lambda^s)^*$ , the functional  $\Lambda \in [L^1(\mathbf{X}, \mathcal{B}(\mathbf{X}), \lambda^s)]^*$  is of the form:

$$\Lambda(f) \equiv \int_{\mathbf{X}} g(x) f(x) dx \quad \text{for some } g \in L^\infty(\mathbf{X}, \mathcal{B}(\mathbf{X}), \lambda^s).$$

**PRELIMINARIES 3** (weak\* topology on  $L^\infty$ ): Endow the space  $L^\infty(\mathbf{X}, \mathcal{B}(\mathbf{X}), \lambda^s)$  with the topology induced by the weak\*-topology on the space  $[L^1(\mathbf{X}, \mathcal{B}(\mathbf{X}), \lambda^s)]^*$ ; see [Rudin \(2005\)](#), p. 67, 68. The new topological space is denoted by  $(L^\infty(\mathbf{X}, \mathcal{B}(\mathbf{X}), \lambda^s), \mathcal{T}^*)$ . Denote convergence in such topology by  $\rightarrow^*$ . Note that, by definition,  $\{g_n\}_{n \in \mathbb{N}} \rightarrow^* g$  if and only if

$$\int_{\mathbf{X}} f(x) g_n(x) dx \rightarrow \int_{\mathbf{X}} f(x) g(x) dx \quad \forall f \in L^1(\mathbf{X}, \mathcal{B}(\mathbf{X}), \lambda^s).$$

Let  $(\mathbf{X}, \Theta, f, \Theta_0)$  be a hypothesis testing problem. Define

$$\mathcal{C}(\alpha-s) \equiv \left\{ \phi \in \mathcal{C} \mid \mathbb{E}_\theta[\phi(X)] - \alpha \equiv \int_{\mathbf{X}} (\phi(x) - \alpha)f(x; \theta) = 0 \quad \forall \theta \in \text{Bd}\Theta_0 \quad \forall \right\}$$

Let  $(L^\infty(\mathbf{X}, \mathcal{B}(\mathbf{X}), \lambda^s), \mathcal{T}^*)$  be the space of essentially bounded functions topologized with the weak\* topology. For any  $A \subset L^\infty(\mathbf{X}, \mathcal{B}(\mathbf{X}))$ , let  $\mathcal{T}_A^*$  denote the subset topology induced by  $\mathcal{T}^*$

**LEMMA 1:** The set  $\mathcal{C}(\alpha-s)$  is compact relative to  $(\mathcal{C}, \mathcal{T}_\mathcal{C}^*)$ .

**PROOF:** The outline of the proof is the following. I show that the set  $\mathcal{C}(\alpha-s)$  is a sequentially closed subset of  $\mathcal{C}$  with the relative weak\* topology. Then I use the Banach-Alaoglu theorem and the topological separability of  $L^1(\mathbf{X}, \mathcal{B}(\mathbf{X}), \lambda^s)$  to establish the compactness of  $\mathcal{C}(\alpha-s)$ .

*(Sequential Closedness)* Take any convergent sequence of tests  $\phi_n \rightarrow^* \phi$  with  $\{\phi_n\}_{n \in \mathbb{N}} \subseteq \mathcal{C}(\alpha-s)$ . I want to show that  $\phi \in \mathcal{C}(\alpha-s)$ . First, I show that  $\phi(x) \in \mathcal{C}$ ; i.e.,  $\phi \in [0, 1]$  for almost every  $x \in \mathbf{X}$ . Suppose not. Then there exists a measurable set  $A \in \mathcal{B}(\mathbf{X})$  with  $\lambda^s(A) > 0$  such that  $\phi(x) > 1$  or  $\phi(x) < 0$  for all  $x \in A$ . Without loss of generality assume  $\phi(x) > 1$ . Since  $\lambda^s$  is  $\sigma$ -finite, there exists a countable collection  $\{E_n\}_{n \in \mathbb{N}}$  such that  $\cup_{n \in \mathbb{N}} E_n = \mathbf{X}$  and  $\lambda^s(E_n) < \infty$  for every  $n$ . Consider the sequence of sets  $\{A \cap E_n\}_{n \in \mathbb{N}}$ . Note that  $0 \leq \lambda^s(A \cap E_n) < \infty$  for all  $n \in \mathbb{N}$ . In addition, there exists  $N \in \mathbb{N}$  for which  $0 < \lambda^s(A \cap E_N)$ , otherwise  $\lambda^s(A) = \lambda^s(\cup_{n=1}^\infty (A \cap E_n)) \leq \sum_{n=1}^\infty \lambda^s(A \cap E_n) = 0$ . Consider the indicator function  $\mathbb{1}_{A \cap E_N}$ . Since  $0 < \lambda^s(A \cap E_N) < \infty$ , the indicator function  $\mathbb{1}_{A \cap E_N} \in L^1(\mathbf{X}, \mathcal{B}(\mathbf{X}), \lambda^s)$ . Note that

$$\lambda^s(A \cap E_N) < \int_{\mathbf{X}} \mathbb{1}_{A \cap E_N}(x) \phi(x) dx = \lim_{n \rightarrow \infty} \int_{\mathbf{X}} \mathbb{1}_{A \cap E_N}(x) \phi_n(x) dx \leq \lambda^s(A \cap E_N).$$

A contradiction. Therefore  $\phi(x) \leq 1$   $\lambda^s$ -almost everywhere in  $\mathbf{X}$ . An analogous argument yields  $\phi(x) \geq 0$   $\lambda^s$ -almost everywhere. Therefore  $\phi \in \mathcal{C}$ . Now, I need to show that  $\phi \in \mathcal{C}(\alpha-s)$ . By assumption, for every  $\theta \in \text{Bd}\Theta_0$   $f(\cdot; \theta)$  is an element of  $L^1(\mathbf{X}, \mathcal{B}(\mathbf{X}), \lambda^s)$ . Consequently,  $f(\cdot, \theta) \in L^1(\mathbf{X}, \mathcal{B}(\mathbf{X}), \lambda^s)$ . Since  $\phi_n \in \mathcal{C}(\alpha-s)$  for every  $n \in \mathbb{N}$  weak\* convergence yields

$$\begin{aligned} 0 &= \lim_{n \rightarrow \infty} \int_{\mathbf{X}} f(x; \theta) (\phi_n(x) - \alpha) dx &= \left( \lim_{n \rightarrow \infty} \int_{\mathbf{X}} f(x; \theta) \phi_n(x) dx \right) - \int_{\mathbf{X}} f(x; \theta) \alpha dx \\ & &= \int_{\mathbf{X}} f(x; \theta) \phi(x) dx - \int_{\mathbf{X}} f(x; \theta) \alpha dx \\ & &= \int_{\mathbf{X}} f(x; \theta) (\phi(x) - \alpha) dx. \end{aligned}$$

So  $\phi \in \mathcal{C}(\alpha-s)$ . This implies  $\mathcal{C}(\alpha-s)$  is sequentially closed in  $\mathcal{C}$  endowed with the weak\* topology.

*(Compactness)* Let

$$V \equiv \left\{ f \in L^1(\mathbf{X}, \mathcal{B}(\mathbf{X}), \lambda^s) : \int_{\mathbf{X}} |f(x)| dx \leq 1 \right\}$$

Note that  $V$  is a neighborhood of the function  $\mathbf{0}$  in the space  $L^1(\mathbf{X}, \mathcal{B}(\mathbf{X}), \lambda^s)$ . Let

$$(A.1) \quad K \equiv \left\{ g \in L^\infty(\mathbf{X}, \mathcal{B}(\mathbf{X}), \lambda^s) : \left| \int_{\mathbf{X}} f(x)g(x) dx \right| \leq 1 \quad \forall f \in V \right\}.$$

Note that  $\mathcal{C}(\alpha-s) \subseteq \mathcal{C} \subseteq K$ , as for any test  $\left| \int_{\mathbf{X}} f(x) \phi(x) dx \right| \leq \int_{\mathbf{X}} |f(x)| |\phi(x)| dx \leq \int_{\mathbf{X}} |f(x)| dx \leq 1$ . By the Banach-Alaoglu Theorem the set  $K$  is compact in the weak\* topology; see [Rudin](#)

(2005), p. 68, Theorem 3.15. Furthermore, the space  $L^1(\mathbf{X}, \mathcal{B}(\mathbf{X}), \lambda^s)$  is topologically separable as  $(\mathbf{X}, \mathcal{B}(\mathbf{X}), \lambda^s)$  is a separable measure space; see exercise 10, Chapter 1 of Stein (2011). Therefore, Theorem 3.16 in Rudin (2005) p. 70 implies that the topological space  $(K, \mathcal{T}_K^*)$  is compact and metrizable. Since every metrizable space is first-countable—consequently, Frechet-Urysohn—the sequential closure of  $C(\alpha-s)$  coincides with its closure. Therefore, the set  $C(\alpha-s)$  is a closed subset of the compact topological space  $(K, \mathcal{T}_K^*)$ . I conclude that  $(\mathcal{C}(\alpha-s), \mathcal{T}_{\mathcal{C}(\alpha-s)}^*)$  is compact and metrizable. That is, the space of  $\alpha$ -similar tests is weak\* compact. *Q.E.D.*

A.2. *Lemma 2: Tests that minimize risk in  $\mathcal{C}(\alpha\text{-s})$  are admissible in the class of all tests*

DESCRIPTION: Let  $(\mathbf{X}, \Theta, f(x; \theta), \Theta_0)$  be a hypothesis testing problem with a product sample space  $(\mathbf{X}_1, \mathbf{X}_2)$ .  $\Theta_0 \neq \emptyset$  is assumed to be a closed set relative to  $(\Theta, \mathcal{T})$  and such that  $\text{Bd}(\Theta_0) = \Theta_0$ .

RELEVANCE OF LEMMA 2: Lemma 2 will be used to establish part i) of Theorem 1.

**LEMMA 2:** Let  $w_1$  denote a full-support probability measures over  $\text{int } \Theta_1$ . Define the *minimum average risk* over the set of  $\alpha$ -similar procedures as:

$$M(w_1) \equiv \arg \min_{\phi \in \mathcal{C}(\alpha\text{-s})} \int_{\text{int } \Theta_1} R(\phi, \theta) dw_1(\theta),$$

and suppose that Assumption F0 holds. Then,

**L2a:** If the sample space  $\mathbf{X}$  is topologically separable:

$$M(w_1) \neq \emptyset.$$

**L2b:** Under Assumption F0,

$$\phi^* \in M(w_1) \implies \phi^* \text{ is admissible in } \mathcal{C}(\alpha\text{-s}).$$

**L2c:** Under Assumption F0:

$$\phi^* \in M(w_1) \implies \phi^* \text{ is admissible in } \mathcal{C}.$$

**Proof of L2a:** For simplicity, assume that  $w_1$  has associated pdf  $p_1$ . I have shown that the class of tests  $\mathcal{C}(\alpha\text{-s})$  is weak\* compact. This class is non-empty, as it contains the randomized test  $\phi(x) = \alpha$ . To establish L2a it will be sufficient to show that the objective function

$$\mathcal{W}^*(\phi) \equiv \int_{\text{int } \Theta_1} R(\phi, \theta) p_1(\theta) d\theta$$

is continuous in the weak\* topology.

**L2a-Step 1** (Fubini's Theorem:) Since the image of any test  $\phi \in \mathcal{C}$  is contained in the interval  $[0, 1]$   $\lambda^s$ -a.e. and  $f(x; \theta) \in L^1(\mathbf{X}, \mathcal{B}(\mathbf{X}), \lambda^s)$  for all  $\theta$ , it follows that  $\left( \int_{\mathbf{X}} \phi(x) f(x; \theta) dx \right) \leq 1$  for every  $\theta \in \Theta$ . Furthermore, since  $p_1(x)$  is also a probability density functions on  $\text{Int}\Theta_1$  and  $\text{Int}\Theta_0$  the following holds

$$\int_{\text{Int}\Theta_1} \left( \int_{\mathbf{X}} \phi(x) f(x; \theta) dx \right) p_1(\theta) d\theta \leq 1 < \infty$$

Therefore, an application of Fubini's theorem in Billingsley (1995), p. 234 yields

$$\int_{\text{int } \Theta_1} R(\phi, \theta) p_1(\theta) d\theta \equiv \int_{\text{int } \Theta_1} \left( \int_{\mathbf{X}} (1 - \phi(x)) f(x; \theta) dx \right) p_1(\theta) d\theta = \int_{\mathbf{X}} (1 - \phi(x)) f_1^*(x) dx$$

where  $f_1^*$  is the "integrated" likelihood given by

$$(A.2) \quad f_1^*(x) \equiv \int_{\text{int } \Theta_1} f(x; \theta) p_1(\theta) d\theta,$$

Note that  $f_1^*$  is an element of  $L^1(\mathbf{X}, \mathcal{B}(\mathbf{X}), \lambda^s)$ . We can re-write

$$(A.3) \quad \mathcal{W}^*(\phi) \equiv \int_{\mathbf{X}} (1 - \phi(x)) f_1^*(x) dx$$

**L2a-Step 2** (Sequential Continuity of  $\mathcal{W}^*$ ): I now show that  $\mathcal{W}^*$  is continuous on the compact metrizable space  $(\mathcal{C}(\alpha\text{-}\mathcal{G}), \mathcal{T}_{\mathcal{C}(\alpha\text{-}s)}^*)$ . It suffices to establish sequential continuity. Take any sequence of tests  $\phi_n \rightarrow^* \phi$ . Since  $f_1^*$  is an element of  $L^1(\mathbf{X}, \mathcal{B}(\mathbf{X}), \lambda_{\mathbf{X}})$ , convergence in the weak\* topology yields

$$\int_{\mathbf{X}} \phi_n(x) f_1^*(x) dx \rightarrow \int_{\mathbf{X}} \phi(x) f_1^*(x) dx.$$

Therefore, the continuity of  $\mathbf{W}$  implies

$$\begin{aligned} \mathcal{W}^*(\phi_n) &\equiv 1 - \int_{\mathbf{X}} \phi_n(x) f_1^*(x) dx \rightarrow 1 - \int_{\mathbf{X}} \phi(x) f_1^*(x) dx, \\ &= \mathcal{W}^*(\phi). \end{aligned}$$

Therefore,  $\mathcal{W}^*$  is a continuous functional defined on the compact space  $(\mathcal{C}(\alpha\text{-}s), \mathcal{T}_{\mathcal{C}(\alpha\text{-}s)}^*)$ , and  $\mathcal{C}(\alpha\text{-}s) \neq \emptyset$ , as it contains the test  $\phi(x) = \alpha$ . This implies  $M(w_1) \neq \emptyset$ .

**L2b** : Let  $\phi^* \in M(w_1)$ . I show that if  $\phi' \in \mathcal{C}(\alpha\text{-}s)$  satisfies

$$(A.4) \quad \mathbb{E}_{\theta}[\phi'(X)] \geq \mathbb{E}_{\theta}[\phi^*(X)] \quad \forall \theta \in \Theta_1$$

then

$$(A.5) \quad \mathbb{E}_{\theta}[\phi'(x)] = \mathbb{E}_{\theta}[\phi^*(x)] \quad \forall \theta \in \Theta_1.$$

Consequently, there is no test  $\phi' \in \mathcal{C}(\alpha\text{-}s)$  that “weakly dominates”  $\phi^*$ ; i.e,  $R(\phi', \theta) \leq R(\phi^*, \theta)$  with strict inequality for some  $\theta$ .

Suppose (A.4) hold, but (A.5) does not. Then, the following is true:

$$C1 \text{ There exists } \tilde{\theta} \in \Theta_1 \text{ such that } \Delta_{\phi', \phi^*}(\tilde{\theta}) \equiv \mathbb{E}_{\tilde{\theta}}[\phi'(X)] - \mathbb{E}_{\tilde{\theta}}[\phi^*(X)] > 0$$

C1 and the continuity of  $\Delta_{\phi', \phi^*}(\cdot)$  at  $\tilde{\theta}$  implies the existence of an open neighborhood  $\tau_{\tilde{\theta}}$  for which  $\Delta_{\phi', \phi^*}(\theta) > 0$  for all  $\theta \in \tau_{\tilde{\theta}}$ . Note that  $\Theta_1 \neq \emptyset$  is an open set. It follows that the set  $\mathcal{S}_{\tilde{\theta}}$  defined by  $\mathcal{S}_{\tilde{\theta}} \equiv \tau_{\tilde{\theta}} \cap \Theta_1$  satisfies three properties: it is non-empty, it is open, and it is contained in  $\Theta_1$ . Since  $w_1(\theta)$  has full support on  $\text{Int}\Theta_1$ ,  $\int_A dw_1(\theta) > 0$  for any open set  $A$  contained in  $\Theta_1$ . Note that  $\Delta_{\phi', \phi^*}(\theta) > 0$  for all  $\theta \in \mathcal{S}_{\tilde{\theta}}$  and (A.4) implies

$$\int_{\Theta_1} \left( \int_{\mathbf{X}} (1 - \phi'(x)) f(x; \theta) dx \right) dw_1(\theta) < \int_{\Theta_1} \left( \int_{\mathbf{X}} (1 - \phi^*(x)) f(x; \theta) dx \right) dw_1(\theta)$$

This contradicts the fact that  $\phi^* \in M(w_1)$ . I conclude C1 cannot hold.

Therefore, (A.4) implies (A.5). I conclude that  $\phi^*$  is admissible in  $\mathcal{C}(\alpha\text{-}s)$ .

**L2c** : I now show that a test  $\phi^* \in M(w_1)$  is admissible in the class of all tests. This proof is based on the arguments provided in [Chernozhukov et al. \(2009\)](#). The proof is divided into two steps.

**STEP 1:** First I show that if  $\phi' \in \mathcal{C}$  satisfies

$$(A.6) \quad \mathbb{E}_\theta[\phi'(X)] \leq \mathbb{E}_\theta[\phi^*(X)] \quad \forall \theta \in \Theta_0$$

and

$$(A.7) \quad \mathbb{E}_\theta[\phi'(X)] \geq \mathbb{E}_\theta[\phi^*(X)] \quad \forall \theta \in \Theta_1$$

with some strict inequality, then  $\phi'$  is  $\alpha$ -similar on  $\text{Bd}\Theta_0 = \Theta_0$ . Consequently, any test  $\phi'$  that “weakly dominates”  $\phi^*$  (i.e,  $R(\phi', \theta) \leq R(\phi^*, \theta)$  with strict inequality for some  $\theta$ ) must be  $\alpha$ -similar on the boundary of  $\Theta_0$ .

Let  $\mathcal{C}_{ns} \subset \mathcal{C}$  be the class of tests that are not similar on the boundary of  $\Theta_0$ . This is,  $\phi \in \mathcal{C}_{ns}$  if and only if there exists  $\theta, \theta' \in \text{Bd}\Theta_0$  such that  $\mathbb{E}_\theta[\phi(x)] \neq \mathbb{E}_{\theta'}[\phi(x)]$ . Partition  $\mathcal{C}$  according to  $\mathcal{C}_{ns}$  so that  $\mathcal{C} \equiv \mathcal{C}_{ns} \cup (\mathcal{C} \setminus \mathcal{C}_{ns})$ . Take any test  $\phi' \in \mathcal{C}_{ns}$  that satisfies (A.6). Since  $\phi'$  is an element of  $\mathcal{C}_{ns}$  and  $\Theta_0$  contains its boundary (as it is closed), there exists  $\theta \in \text{Bd}\Theta_0$  such that  $\Delta_{\phi', \phi^*}(\theta) \equiv \mathbb{E}_\theta[\phi'(X)] - \mathbb{E}_\theta[\phi^*(X)] < 0$ . Because  $\Delta_{\phi', \phi^*}(\theta) < 0$  and the function  $\Delta_{\phi', \phi^*}(\cdot)$  is continuous at  $\theta$ , there exists an open neighborhood  $\tau_\theta \in \mathcal{T}$  such that  $\Delta_{\phi', \phi^*}(\theta) < 0$  for all  $\theta \in \tau_\theta$ . Since  $\theta$  is an element of  $\text{Bd}\Theta_0$ , then  $\tau_\theta \cap \Theta_1 \neq \emptyset$ . The latter implies there exists  $\theta_1 \in \Theta_1$  such that  $\Delta_{\phi', \phi^*}(\theta_1) = \mathbb{E}_{\theta_1}[\phi'(X)] - \mathbb{E}_{\theta_1}[\phi^*(X)] < 0$ . Therefore, equation (A.6) and (A.7) cannot hold. We conclude there is no test  $\phi' \in \mathcal{C}_{ns}$  that satisfies (A.6) and (A.7).

Since  $\mathcal{C}_{ns}$  partitions  $\mathcal{C}$ , a test  $\phi' \in \mathcal{C}$  that satisfies (A.6) and (A.4) must be an element of  $\mathcal{C} \setminus \mathcal{C}_{ns}$  (as  $\phi' \notin \mathcal{C}_{ns}$ ). Equation (A.6) implies  $\phi'$  is  $\alpha'$ -similar on the boundary with  $\alpha' \leq \alpha$ . Two cases follow:  $\alpha' < \alpha$  or  $\alpha' = \alpha$ . In the first case, the argument in the previous paragraph implies that  $\phi'$  will violate (A.4). We conclude that any test that satisfies (A.6) and (A.4) must be  $\alpha$ -similar on  $\text{Bd}\Theta_0 = \Theta_0$ .

**STEP 2:** Since  $\phi^* \in M(w_1)$ ,  $\phi^*$  is admissible in  $\mathcal{C}(\alpha-s)$ . Therefore, there is no  $\alpha$ -similar-on-the-boundary test such that  $R(\phi', \theta) \leq R(\phi^*, \theta)$  with strict inequality for some  $\theta \in \Theta$ . Since —by Step 1— any test  $\phi' \in \mathcal{C}$  that satisfies (A.6) and (A.4) must be  $\alpha$ -similar on  $\text{Bd}\Theta_0$ , I conclude  $\phi^*$  is admissible in  $\mathcal{C}$

## A.3. Proof of Theorem 1

PROOF OF PART I): The proof of the first part follows directly from Lemma 2. Let  $\mathcal{C}(\alpha - s)$  denote the class of  $\alpha$ -similar tests. Simply note that:

$$\begin{aligned} \text{WAP}(\phi_{\text{WAP}}^{w,\alpha}, w) &\equiv \int_{\Theta_1} \left( \int_{\mathbf{X}} \phi_{\text{WAP}}^{w,\alpha}(x) f(x; \theta) dx \right) dw(\theta) \\ &= \int_{\Theta_1} \mathbb{E}_{\theta} [\phi_{\text{WAP}}^{w,\alpha}(x)] dw(\theta) \\ &= 1 - \int_{\Theta_1} R(\phi_{\text{WAP}}^{w,\alpha}, \theta) dw(\theta) \\ &\quad \text{(by the definition of risk function)} \\ &\geq \text{WAP}(\phi, w), \quad \forall \phi \in \mathcal{C}(\alpha - s) \end{aligned}$$

This implies that

$$\int_{\Theta_1} R(\phi_{\text{WAP}}^{w,\alpha}, \theta) dw(\theta) \leq \int_{\Theta_1} R(\phi, \theta) dw(\theta), \quad \forall \phi \in \mathcal{C}(\alpha - s)$$

This means that WAP-similar tests of level  $\alpha$  are average risk minimizers subject to an  $\alpha$ -similarity constraint. Since  $\Theta_0$  is assumed open, then  $\text{int}\Theta_1 = \Theta_1$ . Lemma 2, part c, implies that  $\phi_{\text{WAP}}^{w,\alpha}$  is admissible in the class of all tests.

*Q.E.D.*

PROOF OF PART II): The proof is based on the essentially complete class theorem. See Theorem 2.9.2 and 2.10.3 in [Ferguson \(1967\)](#). See also [Le Cam \(1986\)](#), Chapter 2, Theorem 1.

Note first that the class  $\mathcal{C}(\alpha - s)$  is essentially complete relative to itself (as it contains all the  $\alpha$ -similar tests). Note that the set  $\mathcal{C}(\alpha - s)$  is weak\* compact by Lemma 1. In addition, the risk function of the testing problem  $R(\phi, \theta)$  is—by definition of weak\* topology—continuous (in  $\phi$ ) for all  $\theta \in \Theta$ . This verifies the assumptions of Theorem 2.9.2, p. 85, in [Ferguson \(1967\)](#).

Following Definition 3 [Ferguson \(1967\)](#) p. 50,  $\phi^* \in \mathcal{C}(\alpha - s)$  is said to be an extended Bayes test if for every  $\epsilon > 0$  there is a prior distribution  $w_{\epsilon}(\theta)$  such that:

$$\int_{\Theta_1} R(\phi^*, \theta) dw_{\epsilon}(\theta) \leq \int_{\Theta_1} R(\phi_{\text{WAP}}^{w,\alpha}, \theta) dw_{\epsilon}(\theta) + \epsilon.$$

Theorem 2.10.3 in [Ferguson \(1967\)](#) p. 87 implies that the set of *extended Bayes tests* in  $\mathcal{C}(\alpha - s)$  is *essentially complete*. This essential completeness means that for any other test  $\phi \in \mathcal{C}$  there is a test  $\phi^*$  extended Bayes in  $\mathcal{C}(\alpha - s)$  such that:

$$R(\phi^*, \theta) \leq R(\phi, \theta)$$

for all  $\theta$ . Since  $\phi$  is admissible and  $\alpha$ -similar  $R(\phi^*, \theta) \leq R(\phi, \theta)$  for all  $\theta$  implies that  $R(\phi^*, \theta) = R(\phi, \theta)$ . Therefore, any admissible,  $\alpha$ -similar test is risk equivalent to an extended Bayes test. This implies that for any  $\epsilon > 0$  there is a probability measure  $w_{\epsilon}$  such that

$$\text{WAP}(\phi, w_{\epsilon}) = \text{WAP}(\phi^*, w_{\epsilon}) \geq \text{WAP}(\phi_{\text{WAP}}^{w_{\epsilon}, \alpha}, w_{\epsilon}) - \epsilon$$

Consequently, any admissible,  $\alpha$ -similar test is an extended WAP-similar test of level  $\alpha$ .

*Q.E.D.*



A.4. *Lemma 3: WAP-similar tests with a boundedly complete, null-sufficient statistic.*

PRELIMINARIES: This section generalizes a well-known observation in the IV literature: maximizing constrained average power is straightforward whenever there is a boundedly-complete, null-sufficient statistic. Consider the following assumptions.

**ASSUMPTION F1** (NULL SUFFICIENCY): There is a partition of the data  $X = (x_1, x_2)$  such that the conditional density of  $x_1$  given  $x_2$  in the statistical model  $f(x_1, x_2; \theta)$  satisfies:

$$(A.8) \quad f(x_1|x_2; \beta_0) \equiv f(x_1|x_2; \theta) = f(x_1|x_2; \theta') \quad \forall \theta, \theta' \in \Theta_0.$$

The statistic  $x_2$  arising from such partition of the data will be called a *null-sufficient statistic*.

It is well known that a null-sufficient statistic can be used to control the *null* rejection probability of a test in a two-sided problem with a nuisance parameter [Ferguson (1967), Moreira (2003), Andrews et al. (2006), Lehmann and Romano (2005)].

Let  $h(x_2; \theta)$  denote the marginal density of the null-sufficient statistic  $x_2$  based on the statistical model  $f(x_1, x_2; \beta, \pi)$ .

**ASSUMPTION F2:** (BOUNDED COMPLETENESS): For any bounded measurable function  $m : \mathbf{X}_2 \rightarrow \mathbb{R}$ , the marginal densities of the null sufficient statistic are such that:

$$\int m(x_2)h(x_2; \theta)dx_2 = 0, \quad \forall \theta \in \Theta_0 \implies m(x_2) = 0,$$

except, perhaps, in a set that has zero measure under every element of the collection  $\{h(\cdot, \theta)\}_{\theta \in \Theta_0}$ .<sup>14</sup>

Theorem 4.3.1 in Lehmann and Romano (2005) provides a sufficient condition to guarantee that a family of distributions is complete, and thus, boundedly complete. In the IV example studied in this paper, it will be sufficient to show that the set  $\Theta_0$  contains a rectangle of the same dimension as the null-sufficient statistic.

Bounded completeness will be used to show that all similar tests must be “conditionally” similar. This is a well-known result in the theory of statistical hypothesis testing. See Theorem 4.3.2 in Lehmann and Romano (2005).

DESCRIPTION: Lemma 3 will show that under assumptions F1, F2 the test that rejects whenever

$$(A.9) \quad \phi^*(x_1, x_2) \equiv f_{w_1}^*(x_1, x_2)/f(x_1|x_2; \beta_0) > c(x_2; \alpha),$$

is an element of

$$M(w_1) \equiv \arg \min_{\phi \in \mathcal{C}(\alpha-s)} \int_{\text{Int } \Theta_1} R(\phi, \theta) dw_1(\theta)$$

provided  $c(x_2; \alpha)$  is the  $1 - \alpha$  quantile of  $z(X_1, x_2)$  with  $X_1 \sim f(x_1|x_2; \beta_0)$ . This is a well-known result and we reproduce it for the sake of completeness.

RELEVANCE OF LEMMA 3: Lemma 3 implies that the tests in (A.9) are constrained weighted average power maximizers. This property has been discussed in Andrews, Moreira, and Stock (2004), Chernozhukov et al. (2009). Lemma 3 combined with Lemma 2c implies that the tests in (A.9) are

<sup>14</sup>See Lehmann and Romano (2005) p. 115 for the definition of bounded completeness.

admissible in the class of all tests.

**LEMMA 3:** Let  $\phi^*$  be defined as in (A.9) and let  $c(\cdot; \alpha)$  be measurable. Under Assumptions F1, F2  $\phi^* \in M(w_1)$ ; that is,  $\phi^*$  minimizes average risk inside the class of  $\alpha$ -similar tests.

PROOF: Throughout this proof we assume that  $\mathbf{X} = \mathbf{X}_1 \times \mathbf{X}_2$ . Fubini's theorem (L2a-Step 1) and Theorem 4.3.2 in [Lehmann and Casella \(1998\)](#) implies that  $\phi^* \in M(w_1)$  if and only if  $\phi^*$  solves the problem:

$$\begin{aligned} \min_{\phi \in \mathcal{C}} \int_{\mathbf{X}} (1 - \phi(x)) f_1^*(x) dx \\ \int_{\mathbf{X}_1(x_2)} \phi(x_1, x_2) f(x_1|x_2) dx_1 = \alpha \end{aligned}$$

except, perhaps, for  $x_2$  that belong to a set of measure zero under the marginal density of  $h(x_2, \theta)$  for all  $\theta \in \Theta_0$ . Re-write the objective function as

$$\max_{\phi \in \mathcal{C}} \int_{\mathbf{X}} \phi(x) f_1^*(x) dx.$$

The product structure of  $\mathbf{X}$  and the linearity of the integral allows a further expansion of the previous equation:

$$\max_{\phi \in \mathcal{C}} \int_{\mathbf{X}_2} \left( \int_{\mathbf{X}_1} \phi(x_1, x_2) f_1^*(x_1, x_2) dx_1 \right) dx_2.$$

Note first that the Neyman Pearson Lemma in [Ferguson \(1967\)](#) p. 204 implies that for a fixed  $x_2$  the WAP test  $\phi^*(x_1, x_2)$  solves the problem

$$\max_{\phi \in \mathcal{C}} \int_{\mathbf{X}_1} \phi(x_1, x_2) f_1^*(x_1, x_2) dx_1$$

subject to

$$\int_{\mathbf{X}_1} \phi(x_1, x_2) f(x_1|x_2) dx_1 = \alpha.$$

except, perhaps, for  $x_2$  that belong to a set of measure zero under every  $h(x_2, \theta)$ ,  $\theta \in \text{Bd}\Theta_0$ . Hence, to show that  $\phi^*(x_1, x_2) \in M(w_1)$  it only remains to prove that  $\phi^*(x_1, x_2)$  is measurable. That is,  $\phi^*(x_1, x_2) \in \mathcal{C}(\alpha\text{-s})$ . Assumption F0 implies that  $\phi^*(x_1, x_2)$  is continuous in  $x_1$ , for every  $x_2$ . Furthermore, since  $c(\cdot, \alpha)$  is measurable, then  $\phi^*(x_1, x_2)$  is measurable in  $x_2$ , for every  $x_1$ . Therefore,  $\phi^*(x_1, x_2)$  is a Carathéodory function, as defined in [Aliprantis and Border \(2006\)](#), p. 153. Since the sample space  $\mathbf{X}$  is separable (by assumption) and metrizable (for it is a subset of a euclidean space), Lemma 4.5.1 in [Aliprantis and Border \(2006\)](#) p. 153 implies  $\phi^* : \mathbf{X} \rightarrow [0, 1]$  is measurable. *Q.E.D.*

#### A.5. Continuity of the critical value function

Measurability is required for the proof of Theorem 1. This subsection provides two sufficient conditions that imply the continuity of  $c(\cdot; \alpha)$  (and hence, its measurability).

Let  $f(x; \theta)$  denote the statistical model. Consider the following auxiliary assumptions:

**ASSUMPTION F3:** There exists a function  $g(\theta)$  such that:

$$f(x; \theta) \leq g(\theta) \quad \forall x,$$

and  $\int_{\Theta} g(\theta)dw(\theta) < \infty$ .

**ASSUMPTION F4:**  $f(x_1|x_2; \beta_0) > 0$  for every  $(x_1, x_2)$  and  $f(x_1|x_2; \beta_0)$  is continuous in  $(x_1, x_2)$ .

Assumptions F0, F3, F4 imply that  $c(x_2; \alpha)$  is continuous.

PROOF: Note first that Assumption F0 implies that  $f_w^*(x)$  is sequentially continuous in  $x$ . To see this, consider any sequence  $x_n \rightarrow x$ . Assumption F0 i) implies that  $f(x_n; \theta) \rightarrow f(x; \theta)$  for almost every  $\theta \in \Theta$ . Since the weight function  $w(\theta)$  is assumed to satisfy  $f_w^*(x) < \infty$  for every  $x$  then:

$$\begin{aligned} \left| f_w^*(x_n) - f_w^*(x) \right| &= \left| \int_{\Theta} f(x_n; \theta)dw(\theta) - \int_{\Theta} f(x; \theta)dw(\theta) \right| \\ &\leq \int_{\Theta} \left| f(x_n; \theta) - f(x; \theta) \right| dw(\theta). \end{aligned}$$

By Assumption F3, the Dominated Convergence Theorem applies and we can conclude that

$$\int_{\Theta} \left| f(x_n; \theta) - f(x; \theta) \right| dw(\theta) \rightarrow 0.$$

Consequently,  $f_w^*(x_n) \rightarrow f_w^*(x)$ . Furthermore, Assumption F4 implies that the test statistic:

$$z(x_1, x_2) = f_w(x_1, x_2)/f(x_1|x_2; \beta_0)$$

is continuous in  $(x_1, x_2)$ .

Let  $x_{2,n} \rightarrow x_2$  and let  $X_{1,n} \sim f(x_1|x_{2,n}; \beta_0)$ . Consider the sequence of random variables.

$$z(X_{1,n}, x_{2,n}).$$

By Scheffe's theorem and the continuity of  $f(x_1|x_2; \beta_0)$  at  $x_2$ ,  $X_{1,n} \rightarrow X \sim f(x_1|x_2; \beta_0)$ . Therefore, the random vector  $(X_{1,n}, x_{2,n}) \xrightarrow{d} (X, x_2)$ . The continuous mapping theorem implies that

$$z(X_{1,n}, x_{2,n}) \xrightarrow{d} z(X_1, x_2).$$

Lemma 21.2 in [Van der Vaart \(2000\)](#) implies  $c(x_{2,n}; \alpha) \rightarrow c(x_2; \alpha)$  for any sequence  $x_{2,n} \rightarrow x_2$ . Hence, the critical value function is continuous and, consequently, measurable.